

Local toy-model theory with ontic correlated states of composite systems

Berry Groisman

*Centre for Quantum Information and Foundations,
Centre for Mathematical Sciences, University of Cambridge,
Wilberforce Road, Cambridge CB3 0WA, United Kingdom.*

We propose a toy-model theory, that mimics various characteristic features of quantum mechanics. Unlike the toy-models previously studied in the literature, our toy-model allows for an observer to have a full knowledge of a system's real (ontic) state. This is achieved by introducing domains of disjointness, that is by allowing ontic states to be "non-orthogonal". The observer can perform tests which allow her to distinguish between the states in a single domain of disjointness, but not between all ontic states at once. The consequence of this assumption is that the ontic picture is extended to include joint states of two or more systems. This effectively amounts to emergence of entanglement in the model. We argue that these features, albeit being a "non-classical" element in the theory, support the view that quantum-mechanical states are ontic states.

I. EPISTEMIC MODELS VS ONTIC MODELS

Recently, Local Toy Model Theories (LTMT), initially proposed by Hardy [1] and Spekkens [2], have attracted considerable attention. Hardy was motivated by making a case for non-cloning being logically independent of non-locality. Spekkens used his model to support the conjecture that quantum states are states of observer's incomplete knowledge (epistemic states), rather than real states of affairs (ontic states). Following the two original articles few modified models were proposed [3, 4] and the general framework of those models was discussed in various works [5–9]. The main endeavour of all above works was directed at advocating the epistemic view of quantum states [10, 11].

This article makes a case for ontic view of quantum states. We discuss an alternative local toy-model (i.e. model that admits local hidden variable description), where joint states of correlated individual systems exhibit entanglement (thereby reinforcing recent claims that entanglement is logically independent of non-locality [12]).

Our aim is to construct a toy-model theory in which states of physical systems imitate quantum states, yet the model is not a restriction of quantum theory. One of the central features of quantum mechanics is the existence of non-commutative observables and restrictions on their simultaneous measurements (uncertainty principle). Our model must mimic these features. Our main thesis will be that in the process constructing toy-models with those properties we face two options.

First, we can avoid introducing entangled states by postulating restrictions on our knowledge of the system's state of affairs. This is the route taken by Spekkens. The basic premiss here is that underlying ontic reality of a system is exhausted by a number of mutually disjoint ontic states. In order to achieve non-commutativity and no-cloning, one is forced to introduce limits on the observer's access to this underlying ontic reality, thereby giving (inevitable) rise to a distinction between ontic states and epistemic states - states to which the observer had access to. The ontic reality of states is maintained here only on the level of individual systems. Correlated (joint) states of composite systems are mixtures of ontic states and do not have an ontic status of their own.

Second, we allow the observer to have full access to ontic states. However, we must put additional restrictions on those states in order to mimic non-commutativity, complementarity and no-cloning. The reward is, however, the possibility to extend ontic description to joint states, which is effectively amounts to entanglement. Thus, if we allow full knowledge of the system, then entanglement will be the price to pay. The latter option is a good indication that quantum states are ontic states.

This article is organized as follows. In Section II we set up the general framework for our model and introduce the concept of a domain of disjointness. Section III presents the explicit construction of the model starting with two domains of disjointness in Sub-section III A followed by four domains of disjointness in Sub-section III B. This is followed by discussion in Section IV.

II. WHAT IS ONTIC - SINGLE SYSTEM

The main paradigm of the realist's approach to physical reality constitutes the view that a physical system can be in a number of real physical states. In classical physics all possible states of the system are perfectly distinguishable by appropriate tests and measurements. Quantum-mechanical systems exhibit non-orthogonality and complementarity. If our aim is to construct a toy-model that mimics quantum mechanics we must follow one of the following two strategies.

First strategy is based on

Premise I: The basic premise is that underlying ontic reality of any physical system in isolation constitutes of the set of mutually exclusive (disjoint) states. Disjointness of states implies their distinguishability. Therefore, one needs to impose limitations on access to those properties, e.g. limit distinguishability by hand, introduce hidden variables, knowledge balance principle, etc.

Despite the fact that *Premise I* is deeply rooted in our everyday life experience and in classical physics, there is no compelling reason for why this should be taken as a fundamental starting point.

The second strategy is based on the following premise which, albeit being less intuitive, is nevertheless equally valuable and legitimate.

Premise II: The underlying ontic reality of any physical system constitutes of a certain set of states, $|s\rangle \in \{S\}$, which exhausts all possible internal configurations (degrees of freedom) of that system when it is isolated. The states are not necessarily disjoint. There are subsets of mutual disjointness (domains of disjointness) D , each containing only mutually disjoint states. Two states $|s_i\rangle$ and $|s_j\rangle$ are disjoint if and only if they belong to the same domain of disjointness.

It is reasonable to assume that for every states there must be at least one disjoint states, therefore each D must contain at least 2 states. One could raise an immediate objection to Premise II on the basis that the assumption of disjoint ontic states as un-natural. We stress however, that despite the fact that it will be an unusual assumption to make for many physical systems, there is no inherent logical contradiction in this assumption.

Each D corresponds to a test set-up (at least one), M_D , which allows for an observer to distinguish between the states in D with certainty. Such a measurement will not disturb the states in D . A measurement will disturb the states when implemented on the system when it is in a state from a different D and will yield an inconclusive result. We define as measurement update rule in this case: the state of the system after the measurement M_D will be one of the states from D with equal probability. This randomness is inherent property of the system.

Consider two different domains of disjointness D_x and D_y , where the system is in a state $|s\rangle_x \in D_x$, while measurement is M_y . The measurement setting M_y will correspond to a set of disjoint states $D_y = \{|s\rangle_y\}$. In order to provide systematic description of the measurement process we need to provide a rule that relates $|s\rangle_x$ with D_y . Thus, we define a map $c : |s_i\rangle, |s_j\rangle \in D_x \mapsto |s_k\rangle \in D_y$. The action of $c(|s_i\rangle, |s_j\rangle)$ is consistent with the postulate that there is no ontic reality beyond the states in S due to closure of S under c .

We will consider systems with equal number of states in each D . n -level systems is a system with n states in each domain. Let us describe properties of c on the example of $n = 2$.

III. TWO-LEVEL SYSTEMS

A. Two domains of disjointness

Two-level systems correspond to the case of 2 states in each domain of disjointness. First, consider two domains of disjointness $D_x = \{|0\rangle_x, |1\rangle_x\}$ and $D_y = \{|0\rangle_y, |1\rangle_y\}$, where we have four ontic states in total. Each domain is associated with a corresponding two-outcome test, M_x and M_y respectively. The test M_x distinguishes between the states in D_x with certainty. It yields deterministic outcome $m_x = 0; 1$ if prior to the measurement the system was in the state $|0\rangle_x$ or $|1\rangle_x$ respectively. Thus, the states in D_x fully characterize the test set-up (up to the choice of labels of the outcomes).

What happens if M_x is performed when the systems is in one of the states in D_y ? In this model, it yields an inconclusive result. Thus, we define an overlap between the states $|0\rangle_x$ and $|0\rangle_y$, say, as the probability to obtain outcome $m_x = 0$ when the state is $|0\rangle_y$, $p(m_x = 0/0_y) = 1/2$.

The fact, that we can link M_x with the states of D_y implies that we should be able to define a mathematical relation between D_x and D_y . We define it as follows. The states $|0\rangle_x$ and $|1\rangle_x$ can be combined in two different ways, which give $|0\rangle_y$ and $|1\rangle_y$ respectively and vice versa, i.e.

$$c_{+1}(0_x, 1_x) \mapsto 0_y, \quad c_{+1}(0_y, 1_y) \mapsto 0_x \quad (1)$$

$$c_{-1}(0_x, 1_x) \mapsto 1_y, \quad c_{-1}(0_y, 1_y) \mapsto 1_x \quad (2)$$

$c_{\pm 1}$ are maps or rules, which are analogous to coherent superpositions in quantum mechanics and they form an *isomorphism* between D_x and D_y . The role of $c_{\pm 1}$ is two-fold. First, they provide an ontological link between D_x

and D_y . Second, they allow us to give a quantitative prediction of a result of a test M_i performed when a system is in D_j .

Properties of $c_{\pm 1}$: The two maps are rules, according to which the states are combined. Nevertheless, we demand that they obey certain mathematical properties.

Property 1: $c_{\pm 1}$ can be only defined on states from a single D .

Property 2: *Coherent superpositions of a state with itself.*

$$\begin{aligned}
c_{+1}(s, s) &\mapsto s \\
c_{-1}(s, s) &\mapsto \emptyset \\
c_{\pm 1}(s, \emptyset) &\mapsto s \\
c_{\pm 1}(\emptyset, s) &\mapsto (\pm)s \\
c_{+1}(s, -s) &= c_{-1}(s, s) \\
c_{\pm 1}(\emptyset, \emptyset) &\mapsto \emptyset,
\end{aligned} \tag{3}$$

where $\emptyset$ is a null-state. A null-state is not an ontic state of the system. As seen from the forth line each state has its additive inverse. Additive inverse of a state is not ontologically distinct from the state and is not observable. The model would appear as more attractive without additive inverse, however additive inverse is necessary to maintain consistency between the mathematical properties of $c_{\pm 1}$ and permutation transformations (which will be described late-on). The existence of the additive inverse makes our model closely resembling quantum theory in terms of the mathematical structure. Notice also, that c_{+1} does not obey additivity in the sense the arithmetic operation of addition does.

Property 3: Associativity

$$\begin{aligned}
c_{+1}(c_{+1}(s, t), c_{-1}(s, t)) &= c_{+1}(c_{+1}(s, s), c_{-1}(t, t)), \\
c_{-1}(c_{+1}(s, t), c_{-1}(s, t)) &= c_{+1}(c_{+1}(t, t), c_{-1}(s, s)).
\end{aligned} \tag{4}$$

Combination of Properties (3) and (4) provides closure of the set S under $c_{\pm 1}$. Indeed, it yields

$$\begin{aligned}
c_{+1}(c_{+1}(s, t), c_{-1}(s, t)) &= c_{+1}(s, \emptyset) \mapsto s, \\
c_{-1}(c_{+1}(s, t), c_{-1}(s, t)) &= c_{+1}(t, \emptyset) \mapsto t.
\end{aligned} \tag{5}$$

For example,

$$c_{+1}(c_{+1}(|0\rangle_y, |1\rangle_y), c_{-1}(|0\rangle_y, |1\rangle_y)) = c_{+1}(|0\rangle_y, |\emptyset\rangle) \mapsto |0\rangle_y. \tag{6}$$

Since $c_{\pm}$ obeys associativity and distributivity properties, it will be much more convenient to change the notation from $c_{\pm}(|s\rangle, |t\rangle)$ to $|s\rangle \pm_1 |t\rangle$. However, it should be stressed again that despite resembling arithmetic addition and subtraction these are not arithmetic operations.

For example, consider the combination

$$c_{+1}(|0_x\rangle, |1_x\rangle) \equiv |0_x\rangle +_1 |1_x\rangle \mapsto |0_y\rangle. \tag{7}$$

Each of the two superposed states can be represented in turn as a superposition of states from D_y , i.e.

$$[|0_y\rangle +_1 |1_y\rangle] +_1 [|0_y\rangle -_1 |1_y\rangle] \mapsto |0_x\rangle +_1 |1_x\rangle \mapsto |0_y\rangle. \tag{8}$$

Joint states of two systems: We will now extend our description to states of two or more systems. Such a combined state must include at least 16 states, namely

D_{xx}	D_{xz}	D_{zx}	D_{zz}
$ 0\rangle 0\rangle$	$ 0\rangle 0\rangle$	$ 0\rangle 0\rangle$	$ 0\rangle 0\rangle$
$ 0\rangle 1\rangle$	$ 0\rangle 1\rangle$	$ 0\rangle 1\rangle$	$ 0\rangle 1\rangle$
$ 1\rangle 0\rangle$	$ 1\rangle 0\rangle$	$ 1\rangle 0\rangle$	$ 1\rangle 0\rangle$
$ 1\rangle 1\rangle$	$ 1\rangle 1\rangle$	$ 1\rangle 1\rangle$	$ 1\rangle 1\rangle$

Here these 16 states are divided into four domains of disjointness, e.g. $D_{xx} = \{|0\rangle_x|0\rangle_x, |0\rangle_x|1\rangle_x, |1\rangle_x|0\rangle_x, |1\rangle_x|1\rangle_x\}$ and so on. We will exclude sets of the type $\{|0\rangle_x|0\rangle_x, |0\rangle_x|1\rangle_x, |1\rangle_x|0\rangle_z, |1\rangle_x|1\rangle_z\}$ from being classified as domains of disjointness. We postulate that for each two states in a single D the local parts must be either identical or disjoint.

Since *any* two disjoint states can be superposed coherently (postulated), the full set of states will include superposition of states like $|0\rangle_x|0\rangle_x$ and $|1\rangle_x|1\rangle_x$, etc. We will show that no new type of maps is required to superpose joint states. Consider a superposition $c^{(2)}(|0\rangle_x|0\rangle_x, |0\rangle_x|1\rangle_x)$. It is obvious, that only the states of the second system are being actually superimposed while the state of the first system can be factored out, i.e.

$$c_{+1}^{(2)}(|0\rangle_x|0\rangle_x, |0\rangle_x|1\rangle_x) = |0\rangle_x c_{+1}^{(1)}(|0\rangle_x, |1\rangle_x) = |0\rangle_x [|0\rangle_x +_1 |1\rangle_x] \mapsto |0\rangle_x |0\rangle_y. \quad (9)$$

Thus, the map for 2 systems is of the same type as a map for one system and exhibits *distributivity* property. (We will therefore drop the superscript above the map.) In addition, by associativity (??) we obtain a consistent result

$$\begin{aligned} |0\rangle_x|0\rangle_x +_1 |0\rangle_x|1\rangle_x &= |0\rangle_x [|0\rangle_y +_1 |1\rangle_y] +_1 |0\rangle_x [|0\rangle_z -_1 |1\rangle_y] \\ &= [|0\rangle_x|0\rangle_y +_1 |0\rangle_x|1\rangle_y] +_1 [|0\rangle_x|0\rangle_y -_1 |0\rangle_x|1\rangle_y] \\ &= [|0\rangle_x|0\rangle_y +_1 |0\rangle_x|0\rangle_y] +_1 [|0\rangle_x|1\rangle_y -_1 |0\rangle_x|1\rangle_y] \\ &= |0\rangle_x|0\rangle_y +_1 \emptyset = |0\rangle_x|0\rangle_y. \end{aligned} \quad (10)$$

We arrive now at the main issue. Armed with the laws $c_{\pm}$ satisfy we consider superpositions of the states $|0\rangle_x|0\rangle_x$ and $|1\rangle_x|1\rangle_x$:

$$\begin{aligned} |0\rangle_x|0\rangle_x &= [|0\rangle_y +_1 |1\rangle_y][|0\rangle_y +_1 |1\rangle_y] \\ &= |0\rangle_y [|0\rangle_y +_1 |1\rangle_y] +_1 |1\rangle_y [|0\rangle_y +_1 |1\rangle_y] \\ &= [|0\rangle_y|0\rangle_y +_1 |0\rangle_y|1\rangle_y] +_1 [|1\rangle_y|0\rangle_y +_1 |1\rangle_y|1\rangle_y] \\ &= [|0\rangle_y|0\rangle_y +_1 |1\rangle_y|0\rangle_y] +_1 [|0\rangle_y|1\rangle_y +_1 |1\rangle_y|1\rangle_y], \end{aligned} \quad (11)$$

where the second and third equalities are due to distributivity of $c_{\pm 1}$, while the last equality is due to associativity. Similarly,

$$\begin{aligned} |1\rangle_x|1\rangle_x &= [|0\rangle_y -_1 |1\rangle_y][|0\rangle_y -_1 |1\rangle_y] \\ &= |0\rangle_y [|0\rangle_y -_1 |1\rangle_y] -_1 |1\rangle_y [|0\rangle_y -_1 |1\rangle_y] \\ &= [|0\rangle_y|0\rangle_y -_1 |0\rangle_y|1\rangle_y] -_1 [|1\rangle_y|0\rangle_y -_1 |1\rangle_y|1\rangle_y] \\ &= [|0\rangle_y|0\rangle_y -_1 |1\rangle_y|0\rangle_y] -_1 [|0\rangle_y|1\rangle_y -_1 |1\rangle_y|1\rangle_y], \end{aligned} \quad (12)$$

which yields

$$|0\rangle_x|0\rangle_x +_1 |1\rangle_x|1\rangle_x = |0\rangle_y|0\rangle_y +_1 |1\rangle_y|1\rangle_y. \quad (13)$$

Similarly we can calculate other superpositions. Thus, we obtain the following domain of disjointness D_{ii}^c for correlated states, which are obtained by applying maps $c_{\pm 1}$ to product states of individual systems.

$$\begin{aligned} |0\rangle_x|0\rangle_x +_1 |1\rangle_x|1\rangle_x &= |0\rangle_y|0\rangle_y +_1 |1\rangle_y|1\rangle_y \\ |0\rangle_x|0\rangle_x -_1 |1\rangle_x|1\rangle_x &= |0\rangle_y|1\rangle_y +_1 |1\rangle_y|0\rangle_y \\ |0\rangle_x|1\rangle_x +_1 |1\rangle_x|0\rangle_x &= |0\rangle_y|0\rangle_y -_1 |1\rangle_y|1\rangle_y \\ |0\rangle_x|0\rangle_x -_1 |1\rangle_x|0\rangle_x &= |0\rangle_y|1\rangle_y -_1 |1\rangle_y|0\rangle_y \end{aligned} \quad (14)$$

In this domain the states exhibit definite parity with respect to tests performed on the subsystems in the same “basis”. These states are *ontic* states of a combined system. Thus, in our model correlated states of a composite system have ontic status of their own.

The second domain of disjointness for correlated states D_{ij}^c corresponds to definite parity with respect to the tests performed in the complementary “bases”.

$$\begin{aligned} |0\rangle_x|0\rangle_y +_1 |1\rangle_x|1\rangle_y &= |0\rangle_y|0\rangle_x +_1 |1\rangle_y|1\rangle_x \\ |0\rangle_x|0\rangle_y -_1 |1\rangle_x|1\rangle_y &= |0\rangle_y|1\rangle_x +_1 |1\rangle_y|0\rangle_x \\ |0\rangle_x|1\rangle_y +_1 |1\rangle_x|0\rangle_y &= |0\rangle_y|0\rangle_x -_1 |1\rangle_y|1\rangle_x \\ |0\rangle_x|0\rangle_y -_1 |1\rangle_x|0\rangle_y &= |0\rangle_y|1\rangle_x -_1 |1\rangle_y|0\rangle_x \end{aligned} \quad (15)$$

To emphasize the point that coherent superpositions are ontic states on their own right and shorten the notation we introduce the following notation $C_{\pm 1}^{ab} \equiv |0\rangle_a|0\rangle_b \pm |1\rangle_a|1\rangle_b$ and $A_{\pm 1}^{ab} \equiv |0\rangle_a|1\rangle_b \pm |1\rangle_a|0\rangle_b$ (for correlation and anti-correlation respectively). The complete set of ontic states, divided into corresponding domains of disjointness is summarized in the following table.

D_{xx}	D_{yy}	D_{ab}^c	D_{xy}	D_{yx}	D_{ab}^c
$ 0\rangle_x 0\rangle_x$	$ 0\rangle_y 0\rangle_y$	$C_{+1}^{xx} = C_{+1}^{yy}$	$ 0\rangle_x 0\rangle_y$	$ 0\rangle_y 0\rangle_x$	$C_{+1}^{xy} = C_{+1}^{yx}$
$ 0\rangle_x 1\rangle_x$	$ 0\rangle_y 1\rangle_y$	$C_{-1}^{xx} = A_{+1}^{yy}$	$ 0\rangle_x 1\rangle_y$	$ 0\rangle_y 1\rangle_x$	$C_{-1}^{xy} = A_{+1}^{yx}$
$ 1\rangle_x 0\rangle_x$	$ 1\rangle_y 0\rangle_y$	$A_{+1}^{xx} = C_{-1}^{yy}$	$ 1\rangle_x 0\rangle_y$	$ 1\rangle_y 0\rangle_x$	$A_{+1}^{xy} = C_{-1}^{yx}$
$ 1\rangle_x 1\rangle_x$	$ 1\rangle_y 1\rangle_y$	$A_{-1}^{xx} = A_{-1}^{yy}$	$ 1\rangle_x 1\rangle_y$	$ 1\rangle_y 1\rangle_x$	$A_{-1}^{xy} = A_{-1}^{yx}$

The states in each D^c determine two *global* tests, M_{ii}^c and M_{ij}^c , which distinguish between the states within D_{ii}^c and D_{ij}^c respectively. The two test complement the *local* tests, M_x and M_y .

Transformations: For a single system we define 2 types of transformations. First, permutations within a single D , P_x and P_y . Such permutations have an affect on the complementary D , namely the state $|1\rangle$ acquires a minus sign (becomes its own additive inverse), e.g. $P_x|1\rangle_y = -|1\rangle_y$. Second, a transformation between D_x and D_y , $T_{x \leftrightarrow y}$, which is analogous to a rotation of spin in quantum theory.

Dense Coding: This simple model with only two domains of disjointness exhibits the phenomenon of dense coding in a manner completely analogous to QM. This should not surprise us, however, since so far our model looks very similar to a restricted version of quantum mechanics, if it is restricted to $x - z$ plane of the Bloch-sphere with only superpositions of equal amplitudes. The number of domains of disjointness for a single system is not a determining factor, it appears. What is important is the dimensionality of each domain (2), which leads to four (4) correlated coherent states in a D^c and existence of 4 transformations on a single system, which map one correlated state to a itself or the other three. This is exactly how it works in our model.

Consider a state C_{+1}^{xx} shared between Alice and Bob. The four transformations Alice can perform are I , P_x , P_y and $P_y P_x$.

$$\begin{aligned}
I C_{+1}^{xx} &= I C_{+1}^{yy} \mapsto C_{+1}^{xx} = C_{+1}^{yy} \\
P_x C_{+1}^{xx} &= P_x C_{+1}^{yy} \mapsto A_{+1}^{xx} = C_{-1}^{yy} \\
P_y C_{+1}^{xx} &= P_y C_{+1}^{yy} \mapsto C_{-1}^{xx} = A_{+1}^{yy} \\
P_y P_x C_{+1}^{xx} &= P_y P_x C_{+1}^{yy} \mapsto A_{-1}^{xx} = A_{-1}^{yy}
\end{aligned}$$

Teleportation: Consider three systems A_1 , A_2 and B , where systems A_2 and B are in a correlated state C_{+1}^{xx} . System A_1 is in an unknown arbitrary state, i.e. in one of the four ontic states. Consider all possible situations

$$\begin{aligned}
|0\rangle_x C_{+1}^{xx} &= (C_{+1}^{xx} + C_{-1}^{xx})|0\rangle_x + (A_{+1}^{xx} + A_{-1}^{xx})|1\rangle_x \\
|1\rangle_x C_{+1}^{xx} &= (A_{+1}^{xx} - A_{-1}^{xx})|0\rangle_x + (C_{+1}^{xx} - C_{-1}^{xx})|1\rangle_x \\
|0\rangle_y C_{+1}^{xx} &= |0\rangle_y C_{+1}^{yy} = (C_{+1}^{yy} + C_{-1}^{yy})|0\rangle_y + (A_{+1}^{yy} + A_{-1}^{yy})|1\rangle_y = (C_{+1}^{xx} + A_{+1}^{xx})|0\rangle_y + (C_{-1}^{xx} + A_{-1}^{xx})|1\rangle_y \\
|1\rangle_y C_{+1}^{xx} &= |1\rangle_y C_{+1}^{yy} = (A_{+1}^{yy} - A_{-1}^{yy})|0\rangle_y + (C_{+1}^{yy} - C_{-1}^{yy})|1\rangle_y = (C_{-1}^{xx} - A_{-1}^{xx})|0\rangle_y + (C_{+1}^{xx} - A_{+1}^{xx})|1\rangle_y,
\end{aligned} \tag{16}$$

where we used the relation between correlated states written in difference local bases in Eq. (14).

It is clear, that measurement M_{ii}^c , which is analogous to Bell-measurement in quantum theory, will allow the parties to recover the original states at B by applying corresponding corrective transformations on B as follows

$$\begin{aligned}
C_{+1}^{xx} &\rightarrow I \\
C_{-1}^{xx} &\rightarrow P_y \\
A_{+1}^{xx} &\rightarrow P_x \\
A_{-1}^{xx} &\rightarrow P_y P_x,
\end{aligned} \tag{17}$$

where P_x and P_y are permutations in D_x and D_y respectively. Thus, Alice sends 2 bits to Bob, who performs the corresponding correction.




FIG. 1: Two different types of coherent maps (superpositions) connect the three domains of disjointness.

It is tempting to associate a “ball in a box” interpretation of the ontic states, similar to one adopted by Spekkens. Although, it is possible to do for a single system in our model, such an interpretation cannot be extended to joint states due to their ontic nature. Thus, imposing such an interpretation even on a single system would be misleading. The four ontic states $\{|0\rangle_x, |1\rangle_x, |0\rangle_y, |1\rangle_y\}$ exhaust the ontological baggage of the system. There is no more to it. After the observer performs one of the tests M_i , there is nothing she is ignorant about - she has a full knowledge about the system’s state.

One might argue that this model is merely a restriction of quantum theory. This is true. However, we notice the following. First, our main motivation was to construct a model starting with a strong ontic premiss. If it leads to a model which resembles quantum theory, it is already an indication in favour of the ontic view of quantum states. Second, this model serves as an introduction to more sophisticated version which follows.

To summarize, We have considered a simple model with two domains of disjointness. The model has a simple structure: ontic states with coherent maps that relate between different domains of disjointness. There is no complex vector space, no coefficients or weights with which ontic states are superposed. We have shown how the status of an ontic state can be given to a correlated state of two systems.

B. Four domains of disjointness

To achieve a closer analogy with quantum mechanics additional domains of disjointness are required. The natural step would be to add one additional domain, D_z . This additional domain will require an additional type of coherent maps, $c_{\pm 2}$. Thus, for example

$$\begin{aligned} |0\rangle_x + 1 |1\rangle_x &\mapsto |0\rangle_y, & |0\rangle_x + 2 |1\rangle_x &\mapsto |0\rangle_z \\ |0\rangle_x - 1 |1\rangle_x &\mapsto |1\rangle_y, & |0\rangle_x - 2 |1\rangle_x &\mapsto |1\rangle_z \end{aligned} \quad (18)$$

However, if one is trying to pursue this she will soon encounter a basic difficulty - the arrangement becomes frustrated. For example,

$$\begin{aligned} |0\rangle_y + 2 |1\rangle_y &\mapsto |0\rangle_z, & |0\rangle_z + 2 |1\rangle_z &= (|0\rangle_y + 2 |1\rangle_y) + 2 (|0\rangle_y - 2 |1\rangle_y) \mapsto |0\rangle_y \\ |0\rangle_y - 2 |1\rangle_y &\mapsto |1\rangle_z, & |0\rangle_z - 2 |1\rangle_z &= (|0\rangle_y + 2 |1\rangle_y) - 2 (|0\rangle_y - 2 |1\rangle_y) \mapsto |1\rangle_y, \end{aligned} \quad (19)$$

which contradicts Fig. 1. In quantum theory this frustration is resolved by superposing states with complex coefficients. Here we aim to avoid complex vector spaces.

To overcome this difficulty, we will consider four domains of disjointness. Despite the fact, that it takes us away from a direct analogy with QM, it is important to push the limits of the model without engaging with complex amplitudes. We introduce three different types of coherent maps, each connecting a particular D with three other domains as shown below and summarized in Figure 2.

$$\begin{aligned} &|0\rangle_x + 1 |1\rangle_x \mapsto |0\rangle_y, & |0\rangle_y + 2 |1\rangle_y &\mapsto |0\rangle_z, & |0\rangle_z + 1 |1\rangle_z &\mapsto |0\rangle_t, & |0\rangle_t + 2 |1\rangle_t &\mapsto |0\rangle_x, \\ &|0\rangle_x - 1 |1\rangle_x \mapsto |1\rangle_y, & |0\rangle_y - 2 |1\rangle_y &\mapsto |1\rangle_z, & |0\rangle_z - 1 |1\rangle_z &\mapsto |1\rangle_t, & |0\rangle_t - 2 |1\rangle_t &\mapsto |1\rangle_x, \\ \\ &|0\rangle_y + 1 |1\rangle_y \mapsto |0\rangle_x, & |0\rangle_z + 2 |1\rangle_z &\mapsto |0\rangle_y, & |0\rangle_t + 1 |1\rangle_t &\mapsto |0\rangle_z, & |0\rangle_x + 2 |1\rangle_x &\mapsto |0\rangle_t, \\ &|0\rangle_y - 1 |1\rangle_y \mapsto |1\rangle_x, & |0\rangle_z - 2 |1\rangle_z &\mapsto |1\rangle_y, & |0\rangle_t - 1 |1\rangle_t &\mapsto |1\rangle_z, & |0\rangle_x - 2 |1\rangle_x &\mapsto |1\rangle_t, \\ \\ &|0\rangle_x + 3 |1\rangle_x \mapsto |0\rangle_x, & |0\rangle_z + 3 |1\rangle_z &\mapsto |0\rangle_x, & |0\rangle_y + 3 |1\rangle_y &\mapsto |0\rangle_t, & |0\rangle_t + 3 |1\rangle_t &\mapsto |0\rangle_y, \\ &|0\rangle_x - 3 |1\rangle_x \mapsto |1\rangle_x, & |0\rangle_z - 3 |1\rangle_z &\mapsto |1\rangle_x, & |0\rangle_y - 3 |1\rangle_y &\mapsto |1\rangle_t, & |0\rangle_t - 3 |1\rangle_t &\mapsto |0\rangle_y, \end{aligned} \quad (20)$$




FIG. 2: Three different types of coherent maps (superpositions) connect the four domains of disjointness.

We impose the following properties on $c_{\pm 1}$, $c_{\pm 2}$ and $c_{\pm 3}$.

Property 1: Coherent superpositions of states of a single system.

- Maps of the same type combined in the “cascade” way satisfy distributivity and associativity properties as the maps in Section III A. However, $\pm_3$ is not commutative, as will be specified below. Thus, for example,

$$|0\rangle_x +_1 |1\rangle_x = [|0\rangle_y +_1 |1\rangle_y] +_1 [|0\rangle_y -_1 |1\rangle_y] \mapsto |0\rangle_y. \quad (21)$$

- Maps of different types combined obey

$$\begin{aligned} [|0\rangle_i +_2 |1\rangle_i] +_1 [|0\rangle_i -_2 |1\rangle_i] &= [|0\rangle_i +_1 |1\rangle_i] +_2 [|0\rangle_i -_1 |1\rangle_i] = |0\rangle_i +_3 |1\rangle_i \mapsto |0\rangle_j, \\ [|0\rangle_i +_2 |1\rangle_i] -_1 [|0\rangle_i -_2 |1\rangle_i] &= [|0\rangle_i +_1 |1\rangle_i] -_2 [|0\rangle_i -_1 |1\rangle_i] = |0\rangle_i +_3 |1\rangle_i \mapsto |1\rangle_j, \end{aligned} \quad (22)$$

where j is on the diagonal from i in Fig. 2.

- Special properties of $\pm_3$:

$$\begin{aligned} |0\rangle_i +_3 |1\rangle_i &= |1\rangle_i -_3 |0\rangle_i, \\ |0\rangle_i -_3 |1\rangle_i &= |1\rangle_i +_3 |0\rangle_i. \end{aligned} \quad (23)$$

The map $\pm_3$ does not obey commutativity with respect to interchange of the arguments. These unusual properties are designed to satisfy the transformation rules, which will be discussed in the following.

Property 2: Coherent superpositions of joint states.

- Coherent superpositions involving only $\pm_1$ for join states for all D_{ii} with $i = x, y, z, t$ obey distributivity and associativity as in Section III A.
- For all D_{ii} ($i = x, y, z, t$) coherent superpositions where $\pm_1$, $\pm_2$ and $\pm_3$ are mixed obey

$$\begin{aligned} [|0\rangle_i +_2 |1\rangle_i][|0\rangle_i +_2 |1\rangle_i] +_1 [|0\rangle_i -_2 |1\rangle_i][|0\rangle_i -_2 |1\rangle_i] &= |0\rangle_i |0\rangle_i +_2 |1\rangle_i |1\rangle_i, \\ [|0\rangle_i +_2 |1\rangle_i][|0\rangle_i +_2 |1\rangle_i] -_1 [|0\rangle_i -_2 |1\rangle_i][|0\rangle_i -_2 |1\rangle_i] &= |0\rangle_i |1\rangle_i +_2 |1\rangle_i |0\rangle_i, \\ [|0\rangle_i +_2 |1\rangle_i][|0\rangle_i -_2 |1\rangle_i] +_1 [|0\rangle_i -_2 |1\rangle_i][|0\rangle_i +_2 |1\rangle_i] &= |0\rangle_i |0\rangle_i -_2 |1\rangle_i |1\rangle_i, \\ [|0\rangle_i +_2 |1\rangle_i][|0\rangle_i -_2 |1\rangle_i] -_1 [|0\rangle_i -_2 |1\rangle_i][|0\rangle_i +_2 |1\rangle_i] &= |0\rangle_i |1\rangle_i -_2 |1\rangle_i |0\rangle_i, \end{aligned} \quad (24)$$

$$\begin{aligned} [|0\rangle_i +_1 |1\rangle_i][|0\rangle_i +_1 |1\rangle_i] +_2 [|0\rangle_i -_1 |1\rangle_i][|0\rangle_i -_1 |1\rangle_i] &= |0\rangle_i |0\rangle_i +_2 |1\rangle_i |1\rangle_i, \\ [|0\rangle_i +_1 |1\rangle_i][|0\rangle_i +_1 |1\rangle_i] -_2 [|0\rangle_i -_1 |1\rangle_i][|0\rangle_i -_1 |1\rangle_i] &= |0\rangle_i |1\rangle_i +_2 |1\rangle_i |0\rangle_i, \\ [|0\rangle_i +_1 |1\rangle_i][|0\rangle_i -_1 |1\rangle_i] +_2 [|0\rangle_i -_1 |1\rangle_i][|0\rangle_i +_1 |1\rangle_i] &= |0\rangle_i |0\rangle_i -_2 |1\rangle_i |1\rangle_i, \\ [|0\rangle_i +_1 |1\rangle_i][|0\rangle_i -_1 |1\rangle_i] -_2 [|0\rangle_i -_1 |1\rangle_i][|0\rangle_i +_1 |1\rangle_i] &= |0\rangle_i |1\rangle_i -_2 |1\rangle_i |0\rangle_i, \end{aligned} \quad (25)$$

$$\begin{aligned} [|0\rangle_i +_2 |1\rangle_i][|0\rangle_i +_2 |1\rangle_i] +_2 [|0\rangle_i -_2 |1\rangle_i][|0\rangle_i -_2 |1\rangle_i] &= |0\rangle_i |0\rangle_i +_1 |1\rangle_i |1\rangle_i, \\ [|0\rangle_i +_2 |1\rangle_i][|0\rangle_i +_2 |1\rangle_i] -_2 [|0\rangle_i -_2 |1\rangle_i][|0\rangle_i -_2 |1\rangle_i] &= |0\rangle_i |1\rangle_i +_1 |1\rangle_i |0\rangle_i, \\ [|0\rangle_i +_2 |1\rangle_i][|0\rangle_i -_2 |1\rangle_i] +_2 [|0\rangle_i -_2 |1\rangle_i][|0\rangle_i +_2 |1\rangle_i] &= |0\rangle_i |0\rangle_i -_1 |1\rangle_i |1\rangle_i, \\ [|0\rangle_i +_2 |1\rangle_i][|0\rangle_i -_2 |1\rangle_i] -_2 [|0\rangle_i -_2 |1\rangle_i][|0\rangle_i +_2 |1\rangle_i] &= |0\rangle_i |1\rangle_i -_1 |1\rangle_i |0\rangle_i, \end{aligned} \quad (26)$$

$$\begin{aligned}
& [|0\rangle_i +_3 |1\rangle_i][|0\rangle_i +_3 |1\rangle_i] +_1 [|0\rangle_i -_3 |1\rangle_i][|0\rangle_i -_3 |1\rangle_i] = |0\rangle_i |0\rangle_i +_2 |1\rangle_i |1\rangle_i, \\
& [|0\rangle_i +_3 |1\rangle_i][|0\rangle_i +_3 |1\rangle_i] -_1 [|0\rangle_i -_3 |1\rangle_i][|0\rangle_i -_3 |1\rangle_i] = |0\rangle_i |0\rangle_i -_2 |1\rangle_i |1\rangle_i, \\
& [|0\rangle_i +_3 |1\rangle_i][|0\rangle_i -_3 |1\rangle_i] +_1 [|0\rangle_i -_3 |1\rangle_i][|0\rangle_i +_3 |1\rangle_i] = |0\rangle_i |1\rangle_i +_2 |1\rangle_i |0\rangle_i, \\
& [|0\rangle_i +_3 |1\rangle_i][|0\rangle_i -_3 |1\rangle_i] -_1 [|0\rangle_i -_3 |1\rangle_i][|0\rangle_i +_3 |1\rangle_i] = |0\rangle_i |1\rangle_i -_2 |1\rangle_i |0\rangle_i,
\end{aligned} \tag{27}$$

with same rule as (27) when $\pm_1$ and $\pm_2$ are interchanged.

Properties 1 and 2 provide consistent representation of correlated states in different local “bases”. Using notation introduced in Section III A we can write the first correlated domain of disjointness:

$$\begin{aligned}
C_{+1}^{xx} &= C_{+1}^{yy} = C_{+2}^{zz} = C_{+2}^{tt} \\
C_{-1}^{xx} &= A_{+1}^{yy} = C_{-2}^{zz} = A_{+2}^{tt} \\
A_{+1}^{xx} &= C_{-1}^{yy} = A_{+2}^{zz} = C_{+2}^{tt} \\
A_{-1}^{xx} &= A_{-1}^{yy} = A_{-2}^{zz} = A_{-2}^{tt}
\end{aligned} \tag{28}$$

Thus, introducing the forth domain of disjointness allows our model to exhibit properties of correlations which are quite different from those in quantum theory, namely for a single joint state measured in different bases the toy-model exhibits even number of both correlations and anti-correlations. Unlike quantum mechanics, where for three domains of disjointness the number of anti-correlations is always odd, in this model it can be either even or odd. Thus we conjecture, that our model is not a straightforward restriction of quantum theory. The interesting question to be addressed is whether such correlations can be mimicked within quantum theory.

Transformations: In the case of four domains of disjointness we define 3 types of transformations for a single system. First, P_{xz} and P_{yt} permute the states in two domains. When acting on the complementary domains these transformations cause $|1\rangle$ to acquire a minus sign, e.g. $P_{xz}|1\rangle_y = -|1\rangle_y$. It is impossible to permute states in one domain of disjointness only.

Now it is clear that the properties (23) are the consequence of introduction of P_{xz} and P_{yt} .

Dense Coding: Dense coding works in a similar way to two domains of disjointness. Consider the state C_{+1}^{xx} shared between Alice and Bob. It is easy to check using the properties of these state, that Alice can encode four bits of information by acting locally on her system with one of the transformations I , P_{xz} , P_{yt} , or $P_{yt}P_{xz}$.

Teleportation: Consider three systems A_1 , A_2 and B, where systems A_2 and B are in a correlated state C_{+1}^{xx} . System A_1 is in an unknown arbitrary state, i.e. in one of the six possible ontic states. Consider all possible situations

$$\begin{aligned}
|0\rangle_x C_{+1}^{xx} &= (C_{+1}^{xx} +_1 C_{-1}^{xx})|0\rangle_x +_1 (A_{+1}^{xx} +_1 A_{-1}^{xx})|1\rangle_x \\
|1\rangle_x C_{+1}^{xx} &= (A_{+1}^{xx} -_1 A_{-1}^{xx})|0\rangle_x +_1 (C_{+1}^{xx} -_1 C_{-1}^{xx})|1\rangle_x \\
|0\rangle_y C_{+1}^{xx} &= |0\rangle_y C_{+1}^{yy} = (C_{+1}^{yy} +_1 C_{-1}^{yy})|0\rangle_y +_1 (A_{+1}^{yy} +_1 A_{-1}^{yy})|1\rangle_y \\
|1\rangle_y C_{+1}^{xx} &= |1\rangle_y C_{+1}^{yy} = (A_{+1}^{yy} -_1 A_{-1}^{yy})|0\rangle_y +_1 (C_{+1}^{yy} -_1 C_{-1}^{yy})|1\rangle_y \\
|0\rangle_z C_{+1}^{xx} &= |0\rangle_z C_{+2}^{zz} = (C_{+2}^{zz} +_2 C_{-2}^{zz})|0\rangle_z +_2 (A_{+2}^{zz} +_2 A_{-2}^{zz})|1\rangle_z \\
|1\rangle_z C_{+1}^{xx} &= |1\rangle_z C_{+2}^{zz} = (A_{+2}^{zz} -_2 A_{-2}^{zz})|0\rangle_z +_2 (C_{+2}^{zz} -_2 C_{-2}^{zz})|1\rangle_z \\
|0\rangle_t C_{+1}^{xx} &= |0\rangle_t C_{+2}^{tt} = (C_{+2}^{tt} +_2 C_{-2}^{tt})|0\rangle_t +_2 (A_{+2}^{tt} +_2 A_{-2}^{tt})|1\rangle_t \\
|1\rangle_t C_{+1}^{xx} &= |1\rangle_t C_{+2}^{tt} = (A_{+2}^{tt} -_2 A_{-2}^{tt})|0\rangle_t +_2 (C_{+2}^{tt} -_2 C_{-2}^{tt})|1\rangle_t
\end{aligned} \tag{29}$$

The domain of disjointness (28) defines the “Bell test” M_{ii}^c . We use (28) to associate each outcome in terms of different bases in (29) with the corresponding outcomes in terms of the original basis xx :

A_1	C_{+1}^{xx}	C_{-1}^{xx}	A_{+1}^{xx}	A_{-1}^{xx}
$ 0\rangle_x$	$ 0\rangle_x$	$ 0\rangle_x$	$ 1\rangle_x$	$ 1\rangle_x$
$ 1\rangle_x$	$ 1\rangle_x$	$ 1\rangle_x$	$ 0\rangle_x$	$ 0\rangle_x$
$ 0\rangle_y$	$ 0\rangle_y$	$ 1\rangle_y$	$ 0\rangle_y$	$ 1\rangle_y$
$ 1\rangle_y$	$ 1\rangle_y$	$ 0\rangle_y$	$ 1\rangle_y$	$ 0\rangle_y$
$ 0\rangle_z$	$ 0\rangle_z$	$ 0\rangle_z$	$ 1\rangle_z$	$ 1\rangle_z$
$ 1\rangle_z$	$ 1\rangle_z$	$ 1\rangle_z$	$ 0\rangle_z$	$ 0\rangle_z$
$ 0\rangle_t$	$ 0\rangle_t$	$ 1\rangle_t$	$ 0\rangle_t$	$ 1\rangle_t$
$ 1\rangle_t$	$ 1\rangle_t$	$ 0\rangle_t$	$ 1\rangle_t$	$ 0\rangle_t$

It is clear that after receiving the result of the test the second party can complete the protocol by implementing local corrections as follows

$$\begin{aligned}
C_{+1}^{xx} &\rightarrow I \\
C_{-1}^{xx} &\rightarrow P_{yt} \\
A_{+1}^{xx} &\rightarrow P_{xz} \\
A_{-1}^{xx} &\rightarrow P_{yt}P_{xz},
\end{aligned} \tag{30}$$

The protocol utilized one pair of correlated states and 2 bits of classical communication. Our model resembles quantum mechanics in having the number of classical bits sent larger than the amount of bits of classical information needed to specify an ontic state of A_1 . In this respect, our model is stronger, than the model of Spekkens, where those two quantities match.

Teleportation and Dense Coding are often seen as the manifestation of the characteristic aspects and features of quantum theory and we succeeded in mimicking them in our model. In the future work we will explore additional phenomena and protocols.

IV. DISCUSSION

One can commit to the ontic reality of a single physical system in two different ways.

First way is to attribute a set of distinct disjoint physical states to the system. If one follows this route, then in order to mimic quantum features, e.g. complementarity, non-commutativity, no-cloning and so on, it is essential to introduce a limitation on the access to those states. In other words, one must limit an observer's knowledge about the real state of affairs of the system. When constructing a toy-model one needs to introduce such a limitation "by hand", as it was done by Spekkens with the help of *knowledge balance principle*. In this case all transformations and coherent superpositions are defined for the states of the limited knowledge, e.g. for the epistemic states, not the ontic states. This implies a strongly classical ("ball in a box") approach to ontic states, since it is necessary to rest the rules which govern the transformations of epistemic states on the ontic states of a system. Now, when we are trying to extend the model to include multiple systems and introduce correlations, correlated epistemic states will correspond to probabilistic mixture of product ontic states of individual systems. This picture does not admit a *correlated joint ontic states*.

Second way is to relax the requirement from ontic states of a single system and allow them to be non-disjoint. This is the avenue taken in this article. The expectation from all ontic states being mutually disjoint is deeply rooted in our "classical" experience, but is it not justified as a basic for an ontological commitment. In fact, but putting these classical properties in the very foundation of the toy-model theory we condone this theory to be confined to this classical restriction. Thus, the epistemic picture becomes a self-fulfilling prophecy. The benefit from allowing ontic states of a single system to be disjoint are great. It allows to extend the ontic picture to correlated joint states. In the toy-model suggested in this work, correlated joint states are ontic states. Such states cannot be presented as mixtures of product ontic states of individual systems and therefore are non-separable or entangled, using quantum-mechanical terminology. This model might appear less appealing than models based on the epistemic approach. Indeed, one of the attractive features of Spekkens's model is its elegance and simplicity. It is based on very simple plain assumptions about a single system. Our model does not admit such a simple "which box the ball is?"-interpretation and therefore is much less intuitive. However, this model demonstrates that if we attempt to maintain the ontic interpretation on the level of composite system, we have to introduce a mathematical structure which resembles quantum theory. We started with very basic assumptions, but in the process of constructing the toy-model we were forced to attribute to

it mathematical properties very similar to those that govern quantum theory. One might interpret it as a weakness of the model. Nevertheless, it might be also seen as its strength. In other words, if our basic premiss is that the states of individual systems are ontic, but could be disjoint, and that correlated state are ontic as well, the toy-model theory we get resembles quantum theory in term if its mathematical structure. We see it as an evidence in support of the view that quantum states are ontic states. After all, in nature complementarity, non-commutativity and no-cloning could be inherent features of the states of physical systems on the basic (ontological) level, not the level of access, and this possibility should not be discounted.

Acknowledgments

I would like to thank Ian Stewart, Tony Short, Shahar Avin for productive and helpful discussions and Jeremy Butterfield for giving me the opportunity to present the ideas of this work at a one-day conference "Symmetry in Physics", Trinity College, Cambridge, March 2011. I am also grateful to Daniel Burgarth for inviting me to Imperial College to present the earlier version of this work.

-
- [1] L. Hardy, *Disentangling nonlocality and teleportation*, quant-ph/9906123.
 - [2] R.W. Spekkens, *Evidence for the epistemic view of quantum states: A toy theory*, Phys. Rev. A **75**, 032110 (2007).
 - [3] M. Skotiniotis, A. Roy, B.C. Sanders, *On the epistemic view of quantum states*, J. Math. Phys. **49**, 082103 (2008).
 - [4] S.J. van Enk, *A toy model for quantum mechanics*, Found. Phys. **37**, 1447 (2007). In fact, Van Enk's proposed modification makes his model non-local model, that is a model that allows for Bell-inequalities violation. Non-local Toy Models lie outside the scope of this article.
 - [5] S. Friederich, *How to spell out the epistemic conception of quantum states*, arXiv:1101.1975; published in Studies in History and Philosophy of Modern Physics.
 - [6] M. Pasey, *Stabilizer notation for Spekkens' toy theory*, arXiv:1103.5037.
 - [7] G.C. Knee, *Isolation of the Conceptual Ingredients of Quantum Theory by Toy Theory Comparison*, M.Sc. Thesis, <https://workspace.imperial.ac.uk/theoreticalphysics/Public/MSc/Dissertations/2010/George>
 - [8] B. Coecke and B. Edwards, *Toy quantum categories*, arXiv:0808.1037.
 - [9] S. Avin, *A Philosopher's View of the Epistemic Interpretation of Quantum Mechanics*, <http://philsci-archive.pitt.edu/id/eprint/5364>.
 - [10] Schumacher and Westmoreland have proposed Modal quantum theory [11]. The nature of their model places it apart from those of Hardy, Spekkens and others. It seems to us that it cannot be used as a support of either ontic or epistemic view.
 - [11] B. Schumacher and M.D. Westmoreland, *Modal quantum theory*, arXiv:1010.2929[quant-ph]; *Non-contextuality and free will in modal quantum theory*, arXiv:1010.5452[quant-ph].
 - [12] C. Hewitt-Horsman and V. Vedral, *Entanglement without nonlocality*, Phys. Rev. A , **76**, 062319 (2007).